

A Brief History of Space Climatology: From the Big Bang to the Present

Michael Xapsos^{1D}, Senior Member, IEEE

Abstract—Review of space climatology is presented with a view toward spacecraft electronics applications. The origins and abundances of space radiations are discussed and related to their potential effects. Significant historical developments are summarized leading to the inception of space climatology and into the space era. Energetic particle radiation properties and models of galactic cosmic rays, solar energetic and geomagnetic trapped particles are described. This includes current radiation effects issues that models face today.

Index Terms—Big Bang, galactic cosmic rays (GCRs), solar particle events, space climatology, space radiation models, trapped electrons, trapped protons.

I. INTRODUCTION

THIS review is focused on space climatology—the radiation environment observed over an extended period of time for a given location, corresponding to a space mission duration and orbit. Electronic devices and integrated circuits must be designed for this climatology in order to operate reliably. This will be developed by following a timeline starting with the big bang and ending at the present. A description of the early universe from a radiation effects perspective will be presented, featuring the origin and abundances of relevant particles—electrons, protons, neutrons, and heavy ions. An interesting feature here is a recent development that is changing the view of the origin of ultraheavy elements in the Periodic Table. It will be seen that the origin and abundances of radiations are generally related to the effects they cause in electronic devices and even some of the design requirements that are levied. A transitional period leading to modern times will then be discussed that involves the discovery of sunspots, the solar cycle and the sun’s pervasive influence on space climatology. This leads to the main discussion about modern space climatology for galactic cosmic rays (GCRs), solar particle events, and trapped particles. Radiation properties such as elemental composition, fluxes, energies, and dependence on solar cycle phase and spacecraft orbit will be described, with emphasis on variability of these properties. Radiation models used for space system design will be presented along with some current issues and applications. This will bring the

Manuscript received September 27, 2018; revised December 4, 2018; accepted December 4, 2018. Date of publication December 7, 2018; date of current version January 17, 2019. This work was supported by the NASA Living With a Star Space Environment Testbed Program.

The author is with the NASA Goddard Space Flight Center, Greenbelt, MD 20771 USA (e-mail: michael.a.xapsos@nasa.gov).

Color versions of one or more of the figures in this paper are available online at <http://ieeexplore.ieee.org>.

Digital Object Identifier 10.1109/TNS.2018.2885649




Fig. 1. Timeline from the Big Bang to the present [6].

reader up to date and complete the journey along the space climatology timeline.

II. EARLY UNIVERSE

It is now well established that the size of the universe is expanding with time. Therefore, looking backward in time would reveal a universe that encompasses smaller and smaller volumes the farther back we go. Remarkably, scientists have been able to explain many phenomena by continuing to trace this contraction back to a time about 13.8 billion years ago, considered to be the age of the universe. At this point, it is assumed to be a singularity of infinitesimal size and infinitely dense mass. This generally accepted Big Bang Theory of the birth and evolution of the universe is described in a number of interesting publications for a general audience [1]–[5]. Fig. 1 shows an overall timeline beginning with the Big Bang and continuing through different eras to the present [6].

The following discussion of the early universe is limited to the origin and abundances of radiations that are significant for radiation effects in electronic devices and circuits—electrons, protons, neutrons, and heavy ions. It involves three types of nucleosynthesis processes—Big Bang, stellar, and extreme event nucleosyntheses.

A. Big Bang Nucleosynthesis

A tiny fraction of a second after the Big Bang it is theorized that elementary particles called quarks existed. There are six




Fig. 2. Timeline for the first 380000 years after the Big Bang.




Fig. 3. Timeline for the formation of the first stars.

types of quarks—up, down, top, bottom, strange, and charm. The most stable of these are the up and down quarks, which are the building blocks of nucleons. At times on the order of microseconds after the Big Bang, the early universe had expanded and cooled enough to allow quarks to come together and form stable nucleons. Two up quarks and one down quark form a proton while two down and one up quark form a neutron. Electrons, which are known to be particles with no internal structure, also existed a tiny fraction of a second after the Big Bang along with other elementary particles and energy in the form of light. Continued expansion and cooling allowed protons and neutrons to coalesce into simple nuclei. At an age of about 380000 years, the universe had cooled enough to allow electrons to orbit nuclei and form simple atoms, mainly hydrogen and helium. This portion of the timeline is shown in Fig. 2.

B. Stellar Nucleosynthesis

The formation of the elements in the Periodic Table is a complex subject, and there can be more than one pathway to the synthesis of an element. The purpose of Sections II-C and II-D is not to exhaustively describe this for each element but to simply give a general description of elemental origins so they can ultimately be connected to the radiation effects they cause.

Over a long period of time on the order of hundreds of millions of years, the elements created after the Big Bang, primarily hydrogen, began accumulating into gaseous structures such as the iconic image shown in Fig. 3 taken by the Hubble Space Telescope and known as the “Pillars of Creation.”

These features of the Eagle Nebula are about 4 to 5 light years in their largest dimension. A star will be born within these structures when the density of hydrogen atoms is high enough to start fusing. It is believed that this is how the first stars formed.

At this point in time, stars would have consisted almost entirely of hydrogen and helium. The gravitational attraction of the star’s enormous mass is balanced by the energy release of fusion reactions to form helium, and keeps the star from collapsing in on itself. When the hydrogen is mostly used up, the star begins to contract. This raises the temperature of the core and if the star is large enough (much larger than our sun) helium begins to fuse and additional energy is released to balance the gravitational force. Thus, during the lifetime of large stars, a chain of nuclear fusion reactions starting with hydrogen and helium produce elements from carbon to iron in the star’s core. Iron is the element with the highest binding energy in the Periodic Table and is therefore the most stable. When the star’s core is entirely iron, fusion is no longer possible because the reaction requires energy to be provided rather than resulting in its release. The star’s life is then over. It implodes and becomes a supernova as described in Section II-C. This production of the elements from C to Fe was first proposed by Hoyle [3], [7].

C. Extreme Event Nucleosynthesis

There are two basic conditions that are required for the production of ultra-heavy elements, i.e., those heavier than iron. The first is that there must be enormous energy available in order to overcome the unfavorable energetics of forming these ultraheavy elements from lighter elements. The second is that there must be an abundance of neutrons available, which is seen by examining the excess of neutrons relative to protons in the nuclei of the ultraheavy elements in the Periodic Table. There are few known processes in the universe where this could occur. The two most likely happen after the active lifetimes of large stars. One is due to a supernova explosion, which is initiated when a star’s fuel is used up and the core consists entirely of iron. With no remaining energy to support itself against gravity, the star collapses. Protons and electrons are crushed together to form neutrons and there is a tremendous release of energy from the collapse making the production of the ultraheavy elements possible. A second process is the collision/merger of two neutron stars, observed for the first time August 17, 2017 [8]. A neutron star is the remnant of a large star after a supernova explosion that has collapsed to the density of nuclear material and consists mainly of neutrons. Visible light was detected from this event and gave evidence that ultraheavy elements such as platinum and gold were formed in significant amounts. This led some scientists to postulate it could be the dominant process for formation of ultraheavy elements.

D. Abundances and Radiation Effects of the Elements

With that general background on the origin of elements, their abundances are now examined. Fig. 4 presents the solar abundances of elements in the Periodic Table as a function




Fig. 4. Solar abundances of the elements [9].




Fig. 5. Periodic Table of Radiation Effects.

of mass number. This generally represents the elemental abundances of the solar system [9]. Protons and alpha particles existed shortly after the Big Bang so it is not surprising that the elements H and He are the most abundant. The elements ranging from C to Fe are synthesized in stars larger than the sun in nuclear chain reactions. They are therefore less abundant than the lighter elements H and He. Since the sun ejects these heavy elements during solar particle events but cannot synthesize them, this has the interesting consequence that these heavy elements originated in previous generation stars. Finally, note the rapid decline of the elemental abundances beyond Fe. These ultraheavy elements are likely only produced in the rare explosive processes discussed in Section II-C.

A Periodic Table of Radiation Effects can now be constructed that shows different effects these radiations produce. This is shown in Fig. 5 in which the effects are color coded. The blue indicates that the radiation generally produces total dose effects, including both total ionizing dose (TID) and total nonionizing dose (TNID). A green indicates single event effects (SEEs), and a dotted area indicates charging effects. The table is geared toward radiation effects so electrons and neutrons are included alongside protons. The most abundant radiations, electrons, and protons are largely responsible for cumulative total dose effects that require large numbers of particle strikes in devices. The less abundant alpha particles can contribute to total dose effects to a limited extent as

can neutrons. In space, neutrons are produced primarily by interactions of protons with spacecraft materials, planetary atmospheres, and planetary soils. Due to their large numbers, electrons are mainly responsible for charging, another cumulative effect. The heavy elements C through Fe are not abundant enough to contribute significantly to these cumulative effects, but they are important for SEE. Beyond the Fe, Co, Ni group, the elemental abundances and therefore the particle radiation fluxes in space are very low. This is shown in Fig. 5 by shading only a small portion of the elemental box green. It can, however, be important to consider their effects for high confidence level applications such as destructive or critical SEE. The three remaining elements that have not yet been discussed, Li, Be, and B are relatively rare and produced mainly by fragmentation of heavier GCR ions. This will be shown later in Fig. 10.

III. TRANSITION TO MODERN TIMES

Now that the origin of radiations in the early universe has been discussed along with their abundances and effects on electronic devices, let us move on to the transition period to modern times when the era of space climatology emerged. A timeline of this era is shown in Fig. 6.

The telescope was invented in 1608 by the Dutch lens maker Hans Lippershey. Shortly thereafter Galileo Galilei improved its magnification and was the first to use a telescope to study space. These studies could be regarded as the start of modern experimental astronomy. He was one of the first to observe sunspots through a telescope and hypothesized they were part of the solar surface as opposed to objects orbiting the sun.

Today sunspots are regarded as a proxy to solar activity. They are active regions having twisted magnetic fields that inhibit local convection. The region is therefore cooler than its surrounding and appears darker when viewed in visible light. The connection of sunspots to solar activity is seen in Fig. 7, which compares two images taken at the same time, one in visible light and the other in ultraviolet (uv) light. The bright areas in the uv image indicate high activity and correspond almost exactly to the areas of sunspots, as seen in visible light.

Later in the century, in 1687, the first edition of Isaac Newton’s monumental *Principia Mathematica* was published. This historical book mathematically described the laws of motion and the universal law of gravitation. Significantly, it showed that the law of gravitation could be used to derive Kepler’s empirical laws of planetary motion. This could be viewed as the beginning of modern theoretical astronomy.

However, there was something troubling about the orbit of the planet Mercury that could not be entirely explained by Newton’s law of gravitation. In particular, the observed orbital precession did not exactly match the calculations. It was suspected that there may be an unknown planet inside of Mercury’s orbit that was perturbing it and would be difficult to detect due to its proximity to the sun. In 1826, Heinrich Schwabe began a study in an attempt to understand this. It turned out the puzzle of Mercury’s orbit would not be solved until Einstein applied his model of general relativity to it. However, Schwabe became interested in studying sunspots, and 17 years of meticulous studies later he published a




Fig. 6. Timeline for the emergence of space climatology.




Fig. 7. Images taken of the sun at the same time on February 3, 2002. The left image is in visible light and the right image is in ultraviolet light. Credit: ESA and NASA (SOHO).




Fig. 8. Solar cycles 19–24. Credit: WDC-SILSO, Royal Observatory of Belgium.

paper describing the sunspot cycle. The era of modern space climatology began to take form in 1843 with this discovery.

Today it is recognized that understanding the sun's cyclical activity is an important aspect of modeling the space radiation environment. The record of sunspots dates back to the early 1600s, while numbering of sunspot cycles begins in 1749 with cycle number 1. Currently, sunspot cycle 24 is nearly over. The sunspot cycle is approximately 11 years long, but this can vary as can the activity level from one cycle to the next. This 11-year period is often considered to consist of 7 years of solar maximum when activity levels are high and 4 years of solar minimum when activity levels are low. In reality, the transition between solar maximum and solar minimum is a continuous one, but it is sometimes considered to be abrupt for convenience. The last six solar cycles of sunspot numbers are shown in Fig. 8 [10].

Another common indicator of the approximately 11-year periodic solar activity is the solar 10.7-cm radio flux ($F_{10.7}$). This closely tracks the sunspot cycle. The record of $F_{10.7}$ began part way through solar cycle 18 in the year 1947.

The sun's influence on space climatology and space weather is pervasive. It is a source of solar protons and heavy ions, as well as trapped protons and electrons. Furthermore, it modulates these trapped particle fluxes as well as GCR

fluxes entering our solar system. GCR fluxes interact with the atmosphere and are the main source of atmospheric neutrons. These neutrons decay to protons and electrons and supply additional flux to the trapped particle population. The sun is either a source or a modulator of all energetic particle radiations in the near-Earth region. These radiations are discussed in Section IV.

IV. MODERN TIMES—SPACE CLIMATOLOGY

The prior section brings us to the beginning of the modern era of space climatology. It is shown by the timeline in Fig. 9, and marked by the discovery of the energetic space radiations and their impact on electronics that are used in spacecraft.

GCRs were discovered in 1912 by Victor Hess using electroscopes in a balloon experiment at altitudes between 13 000 and 16 000 feet [11]. The penetrating power of this radiation was clear to Hess from these initial observations. It would turn out to be many orders of magnitude more energetic than particles emitted from radioactive materials, which were known at the time. Solar energetic particles were subsequently discovered by Forbush [12] in 1942. It had been known for nearly 100 years prior that bursts of electromagnetic radiation could be emitted by the sun and affect earth communications, but




Fig. 9. Time line for modern space climatology from the year 1900 to the present and its relation to major radiation effects conferences.

this was the first indication that energetic particles could also be a problem. Shortly after that the transistor was invented at Bell Telephone Laboratories in William Shockley's group [13]. The launch of the first satellites, Sputnik I and II, by the Soviet Union in 1957 was followed by the launch of Explorer I and III by the United States in 1958. The Explorer satellites led to the discovery of the Van Allen Belts by James Van Allen [11]. Researchers began to analyze the effects of radiation on bipolar transistors, primarily for United States Department of Defense applications. With the beginning of this paper the first Nuclear and Space Radiation Effects Conference (NSREC) was held at the University of Washington in 1964 [13], [14]. By 1975, SEU was reported to occur in spacecraft [15], although it was apparently observed three years prior to this by the same group for classified work [16]. The NSREC was continuing to expand and held its first Short Course in 1980 [14]. The Radiation and its Effects on Components and Systems (RADECS) Conference began in 1989. By 1991, the NSREC had recognized the importance of space environment research and began to include an environment session in the conference. Twenty-seven more years along the timeline brings us to the most recent NSREC in 2018.

From this perspective, Sections IV-A–IV-D discuss modern space climatology emphasizing the energetic radiations shown in Fig. 9. Section IV-A begins with a definition of space climatology and space weather. Sections IV-B–IV-D discuss properties, models, and current issues for GCRs, solar particle events, and the Van Allen Belts, respectively. Section IV-E then applies the models and shows examples of TID and SEU environments, including the effect of shielding. Depending on which models are used for TID analysis, radiation specifications can be based on either radiation design margin (RDM) or confidence level. These approaches are also reviewed and compared.

A. Definition of Space Climatology and Space Weather

It is not difficult to find long and complex definitions of space climatology and space weather, especially the latter.

TABLE I
CHARACTERISTICS OF GCRS

| Hadron Composition | Energies | Flux | Radiation Effects |
|---|----------------------------|--|-------------------|
| 90% protons 9% alphas 1% heavier ions | Up to $\sim 10^{20}$ eV | 1 to 10 $\text{cm}^{-2}\text{s}^{-1}$ | SEE |

These terms are generally defined here as the condition of the upper atmosphere and beyond, more specifically the conditions of the space radiation environment for a given location or orbit. For space weather, the time period of interest is the short term, e.g., daily conditions, whereas for space climatology, the time period is an extended one such as a mission duration. This has implications for model use in the design and operation of spacecraft. Climatological models are used during the mission concept, planning, and design phases of spacecraft in order to minimize mission risk. These are generally statistical models that allow risk projection well into the future over the mission duration. Space weather models are used during the launch and operation phases in order to manage residual risk. They are generally nowcast or short-term forecast models of the radiation environment. The following discussion deals mainly with the climatological aspects of the radiation environment.

B. Galactic Cosmic Rays

1) *Properties*: GCR are high-energy charged particles that originate outside of our solar system. Some general characteristics are listed in Table I. They are composed mainly of hadrons, the abundances of which are listed in Table I [17]. A more detailed look at the relative abundances compared to solar abundances is shown in Fig. 10. The two abundance distributions are generally similar. The main differences result from fragmentation of GCR ions that tend to smooth out the GCR distribution relative to the solar abundances. This is particularly noticeable for the elements Li, Be, and B ($Z = 3$ to 5), which are produced mainly from fragmentation




Fig. 10. Comparison of the relative abundances of GCR ions (line) and solar system ions (bars). Credit: NASA (<https://imagine.gsfc.nasa.gov/>).




Fig. 11. Differential flux versus energy for GCRs [18].

of heavier GCR ions such as C and O in occasional collisions with interstellar hydrogen or helium. All naturally occurring elements in the Periodic Table (up through uranium) are present in GCR, although there is a steep drop-off for atomic numbers higher than iron ($Z = 26$).

The amazing variation in energy range of GCRs is shown in Fig. 11 based on data compiled by Swordy [18]. Energies can be up to the order of 10^{20} eV, although the acceleration mechanisms to reach such extreme energies are not understood. GCR with energies less than about 10^{15} eV are generally attributed to supernova explosions within the Milky Way galaxy and more recently neutron star collisions. These fluxes, on the order of a few ions $\text{cm}^{-2} \text{s}^{-1}$, are significant for SEE. On the other hand, the origins of GCR with energies

greater than about 10^{15} eV are largely unknown. It is often stated that the origin of GCR with energies beyond 10^{18} eV is extragalactic [19]. A theoretical limit, the Greisen–Zatsepin–Kuzmin limit [20] shown in Fig. 11, is an upper limit in energy that a GCR proton cannot exceed if it travels a long distance as would occur if it originated in another galaxy. The reasoning is that the proton would interact with the omnipresent Cosmic Microwave Background (CMB) and lose energy to it. The CMB is residual electromagnetic radiation left from the Big Bang [4]. However, this limit appears to have been exceeded many times and is a source of controversy. This illustrates how little is known about these ultrahigh energy particles. Fortunately, particle fluxes at these extreme energies are so low that they are not significant for SEE.

2) *Models*: There has been a long-time interest in developing models of GCR fluxes to aid in design of electronic systems, which began with Adams' [21], [22] development of the GCR model in the Cosmic Ray Effects in Microelectronics 1986 (CREME86) code. This section focuses on two popular models used for calculating SEE rates in space, although there are other interesting models that are available [23]–[26].

One model is that developed by Nymmik *et al.* [27] of Moscow State University (MSU). It is currently used in CREME96 [28], the updated version of the 1986 suite of codes hosted on the Vanderbilt University website, <https://creme.isde.vanderbilt.edu>. The other is the Badhwar–O'Neill model developed at the NASA Johnson Space Center [29], [30]. The two models are based on the idea that the energy spectra of GCR ions outside of the heliosphere is given by Local Interstellar Spectra. A diffusion-convection theory of solar modulation is used to describe the GCR penetration into the heliosphere and transport to near earth at 1 Astronomical Unit. This solar modulation is used as a basis to describe the variation of GCR energy spectra over the solar cycle, as shown in Fig. 12 for iron ions [30]. Both models currently use sunspot numbers as input for solar activity leading to solar modulation. The implementation, however, is different. The MSU model uses multiparameter, semiempirical fits to relate the sunspot numbers to GCR intensity. The Badhwar and O'Neill model solves the Fokker–Planck differential equation for the solar modulation parameter as a function of sunspot number. This implementation and various sources of GCR data are described by Xapsos *et al.* [17]. Fig. 13 shows a comparison of the two models with data. Although both of these models are successfully used for SEE applications, the Badhwar–O'Neill model incorporates a broader and more recent database and is used extensively by the medical community.

For SEE analyses, energy spectra such as those shown in Figs. 12 and 13 can be converted to linear energy transfer (LET) spectra. Integral LET spectra for solar maximum and solar minimum conditions are shown in Fig. 14. These spectra include all elements from protons up through uranium. The ordinate gives the flux of particles that have an LET greater than the corresponding value shown on the abscissa. Given the dimensions of the device sensitive volume this allows the flux of particles that deposit a given amount of charge or greater, and therefore an SEE rate, to be calculated in a simple approximation [31]. For some modern devices, however, the




Fig. 12. Illustration of solar modulation for GCR iron ions [30].




Fig. 15. Fluxes for 80 MeV/amu GCR oxygen during solar cycles 1-24 [30].




Fig. 13. Comparison of the MSU [27] and Badhwar-O’Neill 2014 [30] models with data from various sources.




Fig. 14. GCR LET spectra for solar maximum and solar minimum conditions. From CREME96: <https://creme.isde.vanderbilt.edu>.

LET parameter may have shortcomings for calculating SEE rates in space [32].

The LET spectra shown in Fig. 14 are applicable to geosynchronous missions where there is no significant geomagnetic attenuation. The earth’s magnetic field, however, needs to be accounted for at altitudes lower than geosynchronous.

Due to the basic interaction of charged particles with a magnetic field, the particles tend to follow the geomagnetic field lines. Near the equator the field lines tend to be parallel to the earth’s surface. Thus all but the most energetic ions are deflected away. In the polar regions, the field lines tend to point toward or away from the earth’s surface, which allows much deeper penetration of the incident ions. The effect of the geomagnetic field on incident GCR LET spectra can be calculated in CREME96.

3) *Current Issue: Elevated Fluxes during “Deep” and Prolonged Solar Minima:* In Section IV-B2, the solar modulation of GCR flux has been described. Lower solar activity levels result in higher GCR fluxes. As shown in Fig. 8, the most recent complete solar minimum period between cycles 23 and 24, approximately centered at the year 2009, was quite “deep” and prolonged. In fact, it was the deepest solar minimum of the space era and resulted in the highest GCR fluxes observed in this era. This has raised concerns about solar cycles trending toward this behavior and how elevated the GCR fluxes could get in the future [26].

One of the advantages of basing the solar modulation on sunspot numbers is that there is a continuous detailed record of sunspots dating back to 1749. This allows the GCR fluxes to be estimated over this period of time that covers 24 solar cycles. An example is shown in Fig. 15 for 80 MeV/amu oxygen [30]. It is seen that over this extended period of time the peak flux values for each solar minimum have not varied by more than about 30%. The recent deepest minimum of the space era in 2009 can be compared to the deepest since 1750, which occurred in 1810. It can also be compared to the 1977 solar minimum that is used as a default in CREME96, seen to be more of a typical solar minimum. Given this type of variation, the GCR models should be adequate for design of electronic systems as long as appropriate consideration is given to the recent trend in GCR fluxes. The 1977 period could be used for “typical” specifications while the 2009 period could be used for “worst case” specifications.

C. Solar Particle Events

1) *Properties:* Fig. 16 shows schematic showing solar energetic particle production. These particles are likely energized by magnetic reconnection, a process that converts stored




Fig. 16. Solar energetic particle production. Image credits: NASA and ESA.

magnetic energy to kinetic energy, thermal energy, and particle acceleration. Fig. 16 illustrates the difference between the terms solar flare and coronal mass ejection (CME), which are sometimes incorrectly used as being interchangeable. One type of emission process of the sun is electromagnetic in nature. Irradiance is a comparatively low intensity emission that varies with the solar cycle. By contrast, a solar flare is a burst of electromagnetic radiation characterized by a sudden brightening as shown on the right-hand side in Fig. 16. It turns out that solar flares are often, but not always, accompanied by solar energetic particles. The second general type of the sun's emission process is mass emission. The solar wind is a steady stream of plasma (a gas of free ions and electrons) consisting of protons, alpha particles, and electrons in the eV to keV energy range and has an embedded magnetic field. A CME is a large eruption of plasma that carries an embedded magnetic field stronger than that of the solar wind. A CME image is shown on the left-hand side of Fig. 16. A CME that has a high enough speed will drive a shock wave that further accelerates particles. This is analogous to an airplane creating a shock wave if it exceeds the speed of sound. If the CME driven shock reaches earth, it can cause geomagnetic disturbances. CMEs are also a source of solar energetic particles, as shown in Fig. 16. Further properties of solar flares and CMEs are discussed in a review article by Reames [33] giving a detailed account of the many observed differences.

CMEs are the type of solar particle events that are responsible for the major disturbances in interplanetary space and the major geomagnetic disturbances at earth when they impact the magnetosphere. Therefore the focus here is mainly on CMEs. The mass of magnetized plasma ejected in an extreme CME can be on the order of 10^{17} grams. CME speeds can vary from about 50 to 2500 km/s with an average speed of around 450 km/s. It can take anywhere from hours to a few days to reach the earth. Table II lists some further general characteristics of CMEs.

All naturally occurring chemical elements ranging from protons to uranium are present in solar particle events. They can cause permanent damage such as TID and TNID that is

TABLE II
CHARACTERISTICS OF CMEs

| Hadron Composition | Energies | Integral Fluence (>10 MeV/amu) | Peak Flux (>10 MeV/amu) | Radiation Effects |
|--|----------------|---------------------------------------|--|--------------------|
| 96.4% protons 3.5% alphas ~0.1% heavier ions | Up to ~GeV/amu | Up to $\sim 10^{10}$ cm ⁻² | Up to $\sim 10^6$ cm ⁻² s ⁻¹ | TID TNID SEE |




Fig. 17. Differential flux of 25–250 MeV/nucleon C, N, and O measured with IMP-8 spacecraft instrumentation between 1974 and 1996. Superimposed are the sunspot numbers from solar cycles 21 and 22 [34].

due mainly to protons with a small contribution from alpha particles. Heavy ions are not abundant enough to significantly contribute to these cumulative effects. An extreme CME can deposit a few krad(Si) of dose behind 100 mils (2.5 mm) of aluminum shielding. Even though the heavy ion content is a small percentage of the total, it cannot be ignored. Heavy ions, as well as protons and alpha particles in solar particle events, can cause both transient and permanent SEE.

The solar cycle dependence of both solar particle event and GCR fluxes is shown in Fig. 17 in which the

differential flux of all carbon, nitrogen and oxygen ions in the 25 to 250 MeV/nucleon range is shown during the time period 1974–1996 [34]. Superimposed are the sunspot numbers during that time period illustrating the activity of solar cycles 21 and 22. The solar particle event fluxes are seen as the sharp spikes in Fig. 17, which indicate the statistical and periodic nature of these events. Note that the events occur with greater frequency during the solar maximum time periods. They are superimposed on the low-level background flux of GCR approximately on the order of 10^{-4} ($\text{cm}^2 \cdot \text{s} \cdot \text{sr} \cdot \text{MeV/n}$) $^{-1}$ that slowly varies with the solar cycle as discussed in Section IV-B. The GCR fluxes are approximately anticorrelated with the solar cycle.

2) *Models*: There have been a number of climatological models for solar particle events developed over the years for spacecraft design. Due to the stochastic nature of events, confidence level-based approaches have often been used to allow the spacecraft designer to evaluate risk-cost-performance trades for electronic parts [35]. The first such model was King’s [36] analysis of solar cycle 20 data. One “anomalously large” event, the well-known August 1972 event dominated the fluence of this cycle so the model was often used to predict the number of such events expected for a given mission length at a specified confidence level [37]. Using additional data a model from JPL emerged in which Feynman *et al.* [38] showed the distribution of solar proton event magnitudes is continuous between small events and extremely large events such as that of August 1972. The JPL model is a Monte Carlo-based approach [39]. Other probabilistic models followed based on more recent and extensive data. A model from MSU introduced the full solar cycle dependence by assuming the event numbers are directly proportional to sunspot numbers [40]. The NASA Emission of Solar Protons (ESP) and Prediction of Solar Particle Yields for Characterization of Integrated Circuits (PSYCHIC) models are based on Maximum Entropy Theory and Extreme Value Statistics [41], [42]. The European Space Agency (ESA) Solar Accumulated and Peak Proton and Heavy Ion Radiation Environment (SAPPHIRE) model using the Virtual Timelines method invokes a Levy waiting time distribution [43] and continues to evolve [44]. A new model is also under development that updated the database of the ESP model [45] and incorporates a new approach to solar cycle dependence of event numbers [46]. A summary of a number of statistical models is given in [47].

a) *Cumulative fluence models*: Models for cumulative solar proton fluence are useful for evaluating damage due to TID and TNID. They can also be used to determine long-term SEE rates for devices vulnerable to protons. This can be helpful for estimating the probability of a destructive SEE over the course of a mission.

The most straightforward cumulative solar proton fluence model is ESP/PSYCHIC. It is based on measured annual proton fluences during solar maximum. An advantage of this approach is that it is not necessary to know specific details about the time series of events such as the waiting time distribution, for which there are different approaches [43], [48]. It is implicit in the data. This is shown in Fig. 18 where total fluences from 21 solar maximum years are shown as points for




Fig. 18. Cumulative annual solar proton event fluences during solar maximum periods for three solar cycles plotted on lognormal probability paper. The straight lines are fits to the data [42].




Fig. 19. ESP/PSYCHIC model results for cumulative fluence over a 10-year period including 7 years during solar maximum in GEO. Energy spectra are shown for confidence levels ranging from 50 to 99%.

three different energies [42]. This graph is shown on lognormal probability paper on which a lognormal distribution appears as a straight line. The fit distributions can then be used to obtain the lognormal parameters for N -year distributions. An example result is shown in Fig. 19 for 10 years in geostationary earth orbit (GEO). As is the case for all the climatological models discussed above the output spectra are obtained at a user specified level of confidence for the mission duration. The confidence level represents the probability that the calculated spectrum will not be exceeded during the mission.

Comparison of the JPL, ESP/PSYCHIC, and SAPPHIRE models is shown in Fig. 20 for a 2-year solar maximum period at the 95% confidence level [44]. The JPL and SAPPHIRE models are both Monte Carlo-based approaches. It is seen that the largest differences between models occurs at high proton energies. A new statistical model, the Ground Level Enhancement model, is also shown [49]. It is based on randomly sampling parameters from fit proton spectra based on neutron monitor data analyzed by Tylka and Dietrich [50]. This model makes for an interesting comparison because it is




Fig. 20. Comparison of cumulative fluences predicted by solar proton models for 2 years during solar maximum at the 95% confidence level [44].

based on data that are independent of the other models, which are based on space data.

During a space mission the solar particle event fluence that accumulates during the solar maximum time period is often the dominant contribution to the total fluence. A commonly used definition of the solar maximum period is the 7-year period that spans a starting point 2.5 years before and an ending point 4.5 years after a time defined by the maximum sunspot number in the cycle [39]. The remainder of the cycle is considered solar minimum. Fluences that accumulate during solar minimum can be found in a number of publications [40], [43], [51].

Solar heavy ion models are not as advanced as solar proton models primarily because the data are much more limited. A description of uncertainty propagation is given by Truscott *et al.*[52]. For microelectronics applications they are needed to assess SEE. The ESP/PSYCHIC cumulative fluence model for solar heavy ions is described in [53]. Due to the limited data available the probabilistic model is restricted to long-term (approximately 1 year or more) cumulative fluences and not worst case events. The approach taken was to normalize the alpha particle fluxes relative to the proton fluxes based on measurements of the Interplanetary Monitoring Platform-8 (IMP-8) and Geostationary Operational Environmental Satellites (GOES) instrumentation during the time period 1973 to 2001. The energy spectra of major heavy elements—C, N, O, Ne, Mg, Si, S, and Fe—are normalized relative to the alpha particle energy spectra using measurements of the Solar Isotope Spectrometer onboard the Advanced Composition Explorer spacecraft for the 7-year solar maximum period of solar cycle 23. Remaining naturally occurring minor heavy elements in the Periodic Table are determined from measurements made by the International Sun-Earth Explorer-3 spacecraft or an abundance model. Example results for 2 years during solar maximum at the 50% (median) confidence level behind 100 mils of aluminum shielding are shown in Fig. 21.

LET spectra used for SEE analysis have a somewhat unusual shape. Fig. 21 demonstrates that this shape is due to the elemental contributions. Interestingly, this can be related back to




Fig. 21. LET spectra for cumulative fluences of solar protons and heavy ions for two solar maximum years at the 50% confidence level behind 100 mils of aluminum shielding. The total fluence is multiplied by a factor of 1.5 for clarity. Also shown are the contributions to the total LET spectrum due to protons, alphas, $Z = 3$ (Li) to 26 (Fe), and $Z = 27$ to 92 (trans Fe) [53].

the nucleosynthesis of elements in the Periodic Table described previously. The maximum LET that an ion can have in a material is called the Bragg Peak. Therefore on a LET plot such as Fig. 21, the fluence an ion contributes to the total LET spectrum drops sharply to zero at the Bragg Peak. For example, in silicon, this occurs for protons at an LET less than 1. It is seen that protons and alphas produced in Big Bang nucleosynthesis contribute LET values to the total LET spectrum up to about $1 \text{ MeV} \cdot \text{cm}^2/\text{mg}$. Elements formed in stellar nucleosynthesis contribute up to an LET of about $29 \text{ MeV} \cdot \text{cm}^2/\text{mg}$, while elements formed from extreme event nucleosynthesis contribute over the full range of LET values.

b) *Worst case event models:* Another consideration for spacecraft design is the worst case solar particle event that occurs during a mission. It is important to know how high the SEE rate can get during such an event. The most straightforward approach is to design to a well-known large event. The radiation effects community most often uses the October 1989 event while the medical community often uses the August 1972 event. Hypothetical events such as a composite of the February 1956 and August 1972 events have been proposed [54]. There are also event classification schemes in which the magnitudes range from “small” to “extremely large” that may be useful [55], [56]. At one time, the so-called Carrington Event of 1859 was widely quoted as being a worst case event over the last 400 years based on the nitrate record in polar ice cores [57]. However, the glaciology and atmospheric communities disagreed with this interpretation, as the Carrington Event was not observed in most ice cores [58]. Although this event resulted in a severe geomagnetic storm it is now recognized that the solar proton fluences for this event are not reliably known.

The commonly used October 1989 event is provided for use as a worst case scenario in the CREME96 suite of codes at three levels of solar particle event intensity [28]. They are the




Fig. 22. Comparison of a major solar heavy ion event that occurred in November 2001 with the CREME96 “worst day” model. The progression of daily intensities is indicated with the peak intensity occurring on day 2929 of the mission [59]. Note the LET is in units of g^{-1} and values are therefore a factor of 1000 larger compared to other figures in this paper.

“worst week,” “worst day” and “peak flux” models based on proton measurements from the GOES-6 and GOES-7 satellites and heavy ion measurements from the University of Chicago Cosmic Ray Telescope on the IMP-8 satellite. The peak flux model covers the highest 5-min intensity during the event. Comparisons of these models have been made with data taken by the Cosmic Radiation Environment Dosimetry Experiment onboard the Microelectronics and Photonics Test Bed during a very active period of solar cycle 23 [59]. The data show that three major events during this time period approximately equaled the “worst day” model. An example of this is shown by the LET spectra in Fig. 22.

Another approach to worst case event models is to use statistical methods. The idea is analogous to cumulative fluence models where a worst case event would be calculated for a given confidence level and mission duration. There have been several methods proposed for this including extreme value statistics [41], [60], semiempirical approaches [40], and Monte Carlo calculations [43], [44].

The field of extreme value statistics is one with both an extensive theoretical and applied history. It has frequently been used to describe extreme environmental phenomena such as floods, earthquakes, and high wind gusts [61]–[63]. It has turned out to be a useful radiation effects tool when applied to large device arrays such as high density memories [64], gate oxides [65], [66] and sensors [67], [68]. Considering its broad applicability in the radiation effects area, a brief description of the salient features is given here.

Extreme value statistics focuses on the largest or smallest values taken on by a distribution. Thus, the “tails” of the distribution are the most significant. Here, the focus is obtaining the extreme value distribution of a random process when information is known about the initial distribution.

Suppose that a random variable, x , is described by a probability density $p(x)$ and corresponding cumulative distribution $P(x)$. These are referred to as the initial distributions. Fig. 23 shows an initial probability density for a Gaussian distribution [67]. If a number of observations, n , are made of this random variable there will be a largest value within the




Fig. 23. Extreme value distributions for n -values of 10 and 100 compared to the initial Gaussian distribution [67].

n observations. The largest value is also a random variable and therefore has its own probability distribution. This is called the extreme value distribution of largest or maximum values. Examples of these distributions are shown in Fig. 23 for n -values of 10 and 100. Note that as the number of observations increases the extreme value distribution shifts to larger values and becomes more sharply defined. The extreme value distributions can be calculated exactly for any initial distribution. The probability density for maximum values is

$$f_{\max}(x; n) = n[P(x)]^{n-1}p(x). \quad (1)$$

The corresponding cumulative distribution of maximum values is

$$F_{\max}(x; n) = [P(x)]^n. \quad (2)$$

As n becomes large, the exact distribution of extremes may approach a limiting form called the asymptotic extreme value distribution. If the form of the initial distribution is not known but sufficient experimental data are available, the data can be used to derive the asymptotic extreme value distribution. There are three types of asymptotic extreme value distributions of maximum values—the type I or Gumbel, type II, and type III distributions [61]–[63].

With this background, the problem of worst case event models for solar particle events is now considered. In order to determine a worst case event probabilistically, either by extreme value theory or by Monte Carlo simulation, information about the initial distribution must be known. The first description of the complete initial distribution was determined using Maximum Entropy Theory [41]. This is a mathematical procedure for making an optimal selection of a probability distribution when the data are incomplete by avoiding the arbitrary introduction or assumption of information that is not available. It can therefore be argued that this is the best choice that can be made using the available data [69], [70]. The result is a truncated power law in the distribution of event magnitudes, shown in Fig. 24 for the case of >30 -MeV




Fig. 24. Comparison of the truncated power law distribution to three solar cycles of data during solar maximum [41].




Fig. 25. Probability for worst case event proton fluences expected during the indicated time periods during solar maximum [41].

proton event fluences. This describes the essential features of the distribution. The smaller event sizes follow a power law and there is a rapid falloff for very large magnitude events. Note that Fig. 24 also shows the October 1989 event used as a worst case situation in CREME96. A variant of this distribution has subsequently been proposed [71], but there is no significant improvement in the overall fit to data [44], resulting in the use of both functional forms. However, it can be argued that the sharp drop-off for large event sizes shown in the data of reference 44 indicates a truncated power law is more appropriate.

Given the initial distribution of event magnitudes such as the one shown in Fig. 24, the extreme value method can be applied to obtain a worst case event over the course of a mission. However, this situation is a little more complex. The number of events that occur during a mission is variable, so this must be taken into account. If it is assumed the event occurrence is a Poisson process [39], the worst case distribution can be calculated according to [72], [73]. Example results are shown in Fig. 25 for >30-MeV proton event fluences [41].

The probability of exceeding the fluence shown on the y-axis equals one minus the confidence level.

An interesting feature of this model is the “design limit” shown in Fig. 25. A reasonable interpretation is that it is the best value that can be determined for the largest possible event fluence, given limited data. It is not a physical limit but is an objectively determined engineering guideline for use in limiting design costs.

Other worst case statistical models have been developed for both solar proton event fluences and peak fluxes [40], [43], [44], [60], [72]. There are worst case event statistical models for heavy ions, but these are limited due to the lack of data [40], [74]. There is also a probabilistic model for solar electrons that is part of an interplanetary electron model [75].

3) *Current Issue: Use of Statistical Models vs. Worst Case Observations:* As seen in Section IV-C2, there are two types of approaches for evaluating worst case solar particle events. One is to use a worst case observation such as the event that occurred in October 1989, as in CREME96. The other is to use a statistical model to calculate the worst case event that will occur during the mission at a specified level of confidence. Fig. 24 illustrates a set of data that can be used for these approaches. This section compares the approaches and discusses the advantages and disadvantages of each.

The worst case observation approach is straightforward. On the other hand, a statistical model uses an entire database of events and there is much to consider. Events can have very different characteristics in terms of magnitudes (fluence or peak flux), time profile, energy spectra, and heavy ion content. The proton and heavy ion characterization of a worst case observation are self-consistent. This is not necessarily true for the worst case statistical model in which the proton and heavy ion fluxes are analyzed independently. For example, fluxes for different particles can peak at separate times, leaving open different approaches to what characterizes the worst case.

An advantageous feature of the statistical model is that it allows the designer to make risk, cost, performance trades when selecting electronic parts. For example, a higher risk can be assumed in return for a higher performance or less expensive part. By comparison, a worst case observation such as the October 1989 event has little flexibility in the design environment, which is quite severe. This can make requirements difficult to meet for higher risk missions such as CubeSats. Thus, considering the type of mission can be important for deciding on an approach.

Lastly, it is worth noting the current state of development of these models. The worst case observation approach has a long history of successful use. Worst case statistical models for solar protons are also successfully used while heavy ion models are a developing area of research.

D. Van Allen Belts

1) *Trapped Particle Motion in the Magnetosphere:* The earth’s magnetosphere consists of both an external field due to the solar wind and an internal magnetic field. The internal or geomagnetic field originates primarily from within




Fig. 26. Motion of a charged trapped particle in the Earth's magnetic field. After E.G. Stassinopoulos [34].

the earth and is approximately a dipole field. The solar wind and its embedded magnetic field tends to compress the geomagnetic field. During moderate solar wind conditions, the magnetosphere terminates at roughly 10 earth radii on the sunward side. During turbulent magnetic storm conditions, it can be compressed to about six earth radii. The solar wind generally flows around the geomagnetic field and consequently the magnetosphere stretches out to a distance of possibly 1000 earth radii in the direction away from the sun.

The geomagnetic field is approximately dipolar for altitudes up to about four or five earth radii. It turns out that the trapped particle populations are conveniently mapped in terms of the dipole coordinates approximating the geomagnetic field. This dipole coordinate system is not aligned with the earth's geographic coordinate system. The axis of the magnetic dipole field is tilted about 11.5° with respect to the geographic North-South axis and its origin is displaced by a distance of more than 500 km from the earth's geocenter. The standard method is to use McIlwain's (B, L) coordinates [76]. Within this dipole coordinate system, L represents the distance from the origin in the direction of the magnetic equator, expressed in earth radii. One earth radius is 6371 km. B is simply the magnetic field strength. It describes how far away from the magnetic equator a point is along a magnetic field line. B -values are a minimum at the magnetic equator and increase as the magnetic poles are approached. Further background information on the magnetosphere and (B, L) coordinates can be found in [73] and [77].

The basic motion of a trapped charged particle in the geomagnetic field is shown in Fig. 26. Charged particles become trapped because the magnetic field can constrain their motion. The particle spirals around and moves along the magnetic field line. As the particle approaches the polar region, the magnetic field strength increases and causes the spiral to tighten. Eventually, the field strength is sufficient to force the particle to reverse direction. Thus, the particle is reflected between so-called "mirror points" and "conjugate mirror points." In addition, there is a slower longitudinal drift of the path around the earth that is westward for protons and eastward for electrons. This is caused by the radial gradient in the magnetic field. Once a complete azimuthal rotation is made around the earth, the resulting toroidal surface that has been traced out is called a drift shell or L -shell. The L -shell

TABLE III
TRAPPED PROTON CHARACTERISTICS

| L -Shell Values | Energies | Fluxes* (>10 MeV) | Radiation Effects |
|-------------------|------------------|---|--------------------|
| ~ 1 to 10 | Up to \sim GeV | Up to $\sim 10^5$ $\text{cm}^{-2}\text{s}^{-1}$ | TID TNID SEE |

* long-term average




Fig. 27. Trapped proton fluxes >10 MeV mapped in a dipole coordinate system [73].

parameter indicates magnetic equatorial distance from earth's center in number of earth radii and represents the entire drift shell. This provides a convenient global parameterization for a complex population of particles.

2) Trapped Protons:

a) *Properties:* Some of the characteristics of trapped protons and their radiation effects are summarized in Table III and Fig. 27. The L -shell range is from slightly more than 1 at the inner edge of the trapped environment out beyond geosynchronous orbits to an L -value of around 10. The atmosphere limits the belt to altitudes above about 200 km. Trapped proton energies extend up to the GeV range. The energetic trapped proton population with energies >10 MeV is confined to altitudes below 20000 km, while protons with energies of a few MeV or less are observed at geosynchronous altitudes and beyond. The maximum flux of >10 -MeV protons occurs at an L -value around 1.7 and exceeds 10^5 $\text{cm}^{-2}\text{s}^{-1}$. Trapped protons can cause TID, TNID and SEE.

Trapped proton fluxes in low earth orbit (LEO) are approximately anticorrelated with solar cycle activity. This is most pronounced near the belt's inner edge as shown in Fig. 28 [78]. Here $F_{10.7}$, the solar 10.7-cm radio flux, is used as a proxy for solar activity. As solar activity increases the atmosphere expands and causes greater losses of protons to the atmosphere during solar maximum. In addition, there is a decreased production of protons in the atmosphere during solar maximum coming from the Cosmic Ray Albedo Neutron




Fig. 28. Approximate anticorrelation of low altitude trapped proton flux (points) with $F_{10.7}$ as an indicator of solar activity [78].




Fig. 29. Contour plot of proton fluxes >35 MeV in the SAA at an altitude of about 840 km measured by the Polar Orbiting Earth Satellite (POES) from July 1998 to December 2011 [79].

Decay (CRAND) process. The CRAND process is the production of atmospheric neutrons from GCR that subsequently decay to protons (and electrons) and can become trapped. As discussed previously, GCR fluxes are lower during solar maximum.

For spacecraft that have an orbit lower than about 1000 km, the so-called “South Atlantic Anomaly” (SAA) dominates the radiation environment. This anomaly is due to the fact that the earth’s geomagnetic and rotational axes are tilted and shifted relative to each other as discussed before. Thus, part of the proton belt’s inner edge is at lower altitudes in the geographic region around South America. It is shown in Fig. 29 as a contour plot on geographic coordinates for >35 -MeV proton fluxes at an altitude of about 840 km [79].

Higher energy protons are generally fairly stable in the proton belt. However, during the 1990–1991 Combined Release and Radiation Effects Satellite (CRRES) mission the Air Force Research Laboratory (AFRL) discovered the formation of a transient proton belt in the L -shell 2 to 3 region [80]. It is




Fig. 30. Sudden changes in 9.65- to 11.35-MeV trapped proton fluxes caused by solar particle events measured on the Satellite for Scientific Applications (SAC-C) [73].

now known that CMEs can cause geomagnetic storms that suddenly reconfigure the belt. Fig. 30 shows that enhanced fluxes can occur in the L -shell 2 to 3 region if a CME is immediately preceded by another event [73]. Note that although the enhanced flux begins to decay immediately it can remain measurable for well over a year. Fig. 30 also shows that a CME can cause reduction of an enhanced flux. The details of these belt reconfigurations are not fully understood.

b) Models: The general approach to a trapped particle model calculation is to first use an orbit generator to obtain the geographical coordinates of the spacecraft—latitude, longitude, and altitude. Next, the geographical coordinates are transformed to a dipole coordinate system in which the particle population is mapped. The trapped particle environment is then determined external to the spacecraft. The Space Environment Information System (SPENVIS) suite of programs has implemented a number of trapped particle models for unrestricted use at <http://www.spENVIS.oma.be/>.

The well-known Aerospace Proton-8 (AP-8) trapped proton model is the eighth version of a model development effort led by James Vette. Over the years, these empirical models have been indispensable for spacecraft designers and for the radiation effects community in general. The trapped particle models are static maps of the particle population during solar maximum and solar minimum based on data from the 1960s to 1970s. Because these models provide the mean flux values of the environment, a RDM is used for design specifications. Details of the AP-8 model and its predecessors can be found in [81] and [82].

The shortcomings of AP-8 and the need for updates have been discussed [83]. Consequently, there have been a number of notable efforts to develop new trapped proton models [78], [80], [84]–[86]. Comparisons of these models with AP-8 and each other for different orbits are given by Lauenstein and Barth [87].




Fig. 31. Comparison of the AP8 and AP9/IRENE (version 1.5) models for a polar LEO.

Recently, more comprehensive models have been developed. One such model was initially called AP-9 and is now undergoing a name change to the International Radiation Environment Near Earth (IRENE) model [79], [88]. AP9/IRENE allows three methods of calculation. There is a statistical model for the mean or percentile environment. There is a perturbed model that adds measurement uncertainty and data gap filling errors. Third, there is a Monte Carlo capability that includes space weather variations. AP9/IRENE is based on data taken between 1976 and 2016. It does not include solar cycle variation, i.e., output is averaged over the solar cycle. As a result of its probabilistic approach and use of percentiles, confidence levels can be used for design specifications. The other recent comprehensive model is the Global Radiation Earth Environment model (GREEN) [89]. GREEN is an integration of AP-8 with other models that have been developed in order to expand the overall energy and orbital capabilities. Results for the GREEN model were not available at the time of this writing.

Fig. 31 is a comparison of AP-8 and AP-9/IRENE for a polar LEO. The orbital parameters used were those of the Landsat-8 satellite. This provides a reasonable overall comparison as the spacecraft flies through varying portions of the proton belt multiple times each day. Although there are large differences between the models at energies less than 1 MeV, these energies are not significant for most applications. Over most of the remaining energy range, the AP8 model shows higher fluxes during solar minimum compared to solar maximum, as expected, while AP9/IRENE generally results in the highest fluxes. AP9/IRENE also extends to higher energies, which is due to the incorporation of the NASA Van Allen Probes data.

3) *Trapped Electrons*::

a) *Properties*: Some of the characteristics of trapped electrons are summarized in Table IV and shown in Fig. 32. There is both an inner and an outer zone of trapped electrons. These two zones are very different so the characteristics are listed separately. As is also the case for trapped protons the boundaries of the zones are not sharp and they are to some extent dependent on particle energy. For the purposes of this discussion the inner zone is assumed to be between

TABLE IV
TRAPPED ELECTRON CHARACTERISTICS

| | <i>L</i> -Shell Values | Energies | Fluxes* (> 1 MeV) | Radiation Effects |
|------------|------------------------|---------------|---|-------------------------|
| Inner Zone | 1 - 2 | Up to 5 MeV? | uncertain | TID TNID Charging |
| Outer Zone | 3 - 10 | Up to ~10 MeV | Up to ~3x10 ⁶ cm ⁻² s ⁻¹ | |

* long-term average




Fig. 32. Trapped electron fluxes > 1 MeV according to the AE-8 model during solar maximum [73].

L-values of 1 and 2. It was originally thought that electron energies range up to approximately 5 MeV but that has not been observed recently. This electron population tends to remain relatively stable but a long-term average is difficult to ascertain as will be seen in Section IV-C. The outer zone has *L*-values ranging between about 3 and 10 with electron energies generally less than approximately 10 MeV. Here, fluxes peak between *L*-values of 4 and 4.5 and the long-term average for >1 MeV electrons is about 3 × 10⁶ cm⁻² s⁻¹. This zone is very dynamic and the fluxes can vary by orders of magnitude from day to day. An interesting feature of the outer belt is that it extends down to low altitudes at high latitudes. Trapped electrons contribute to TID, TNID and both surface and internal charging effects.

The distribution of trapped particles is a continuous one throughout the inner and outer zones. Between the two zones is a region where the fluxes are at a local minimum during quiet periods. This is known as the slot region. The location of the slot region is assumed to be between *L*-values of 2 and 3 for this discussion. This is an attractive one for certain types of missions due to the increased spatial coverage compared to missions in LEO.

b) *Models*: The long-time standard model for trapped electrons has been the Aerospace Electron-8 (AE-8) model [82], [90]. It consists of two static flux maps of trapped




Fig. 33. Total electron flux before and after a geomagnetic storm compared to a long-term average as measured onboard the UARS [91].

electrons—one for solar maximum and one for solar minimum conditions. Due to the variability of the outer zone electron population, the AE-8 model is valid only for long periods of time. A conservative rule of thumb is that it should not be applied to a period shorter than 6 months.

A feature of the outer zone is its high degree of volatility and dynamic behavior. This results from geomagnetic storms and substorms, which cause major perturbations of the geomagnetic field. Measurements from the Upper Atmosphere Research Satellite (UARS) illustrate the high degree of variability of electron flux levels prior to and after such storms. Fig. 33 shows the electron energy spectra for $3.25 < L \leq 3.5$ after long-term decay from a prior storm (day 235) and two days after a large storm (day 244) compared to the average flux level over a 1000 day period [91]. It is seen for example, at 1 MeV, that the difference in the one-day averaged differential fluxes over a 9-day period is about three orders of magnitude. This illustrates the difference between the long-term average space climate and the short-term space weather in the outer zone.

Due to the volatile nature of the outer zone, it seems natural to resort to probabilistic methods. This is the case for the new AE-9/IRENE trapped electron model [79], [88], which uses the same methodology as described before in the discussion on trapped protons. Other statistical analyses have also been used for both the outer zone and slot region [91]–[94]. Another approach used to describe outer zone fluxes has been to relate them to the level of disturbance of the geomagnetic field by using geomagnetic activity indices such as A_p [95] and K_p [96].

An important orbit in the outer zone that is widely used for telecommunications satellites is GEO. Fig. 34 shows a comparison between the AE8 and AE9/IRENE mean values. AE8 has no solar cycle dependence in GEO so there is no distinction between solar maximum and solar minimum, as was the case in Fig. 31. It is seen that AE8 gives more conservative fluxes over most of the energy range. The group at ONERA, the French National Aerospace Research Center, has also done considerable work on trapped electron models for GEO. Their most recent model is IGE-2006 [97], which gives the option of




Fig. 34. Comparison of the AE8 and AE9/IRENE (version 1.5) models for GEO.

a maximum (worst case), mean or minimum (best case) flux output. When calculation of the mean flux is done in SPENVIS and compared to Fig. 34, results show lower fluxes than both AE8 and AE9/IRENE except at energies approximately less than 0.1 MeV. However, the IGE-2006 model has been incorporated into the group's new comprehensive GREEN model for trapped electrons so more detailed comparisons are deferred until GREEN becomes available for use.

Fig. 35 gives a good overall view of the dynamic behavior of trapped electrons for about a 3.5-year period as measured by Van Allen Probes instrumentation [98]. Fluxes of 0.75-MeV electrons are mapped out according to L -shell values as a function of time. Color coding of electron intensities are shown along the top of the graph. The two boxed areas indicate the most severe storm periods. Fig. 35 shows the volatile nature of the outer zone ($L > 3$). During storm periods electrons can be injected into the slot region ($2 < L < 3$). Here, they are fairly short-lived as the decay period is about 10 days. During severe storms, electrons can also be injected into the inner zone ($1 < L < 2$). Note the stability of the inner zone as the injected electrons decay away very slowly and persist strongly more than a year after the storm.

c) Current issue: the case of the missing electrons: Fig. 35 is a good indicator of the behavior of the electron belts in recent times for energies up to about 0.75 MeV. The inner zone is fairly stable for long periods of time, as evidenced in Fig. 35. When high energy (> 1.5 MeV) electron data are similarly examined as shown in the top portion in Fig. 36 [98], nothing looks out of the ordinary. The outer belt looks volatile and the inner belt appears stable. While inner zone fluxes predicted by models in current use such as AE8 and AE9/IRENE are not large for energies between 1.5 MeV and a maximum of about 5 MeV, they are ordinarily accounted for in radiation effects analysis. However, the top portion in Fig. 36 has not been corrected for background counts, which is mainly due to high energy protons. The Van Allen Probes instrumentation has improved capability in this regard and when background counts are removed the result is shown in the bottom portion of Fig. 36. The high-energy electrons of the inner zone are almost completely gone. In fact, there is no evidence of > 1.5 -MeV




Fig. 35. Fluxes of 0.75-MeV electrons mapped according to *L*-shell as a function of time for approximately 3.5 years. Fluxes are background corrected [98].




Fig. 36. Fluxes of 1.58-MeV electrons mapped according to *L*-shell as a function of time for approximately 3.5 years. The top graph is uncorrected for background counts and the bottom graph is corrected. Note the difference in the inner zone ($1 < L < 2$) [98].

electrons in the inner zone since the Van Allen Probes were launched in 2012. This is the case of the missing electrons.

The question of what happened to this portion of the inner zone remains. Instrumentation prior to the Van Allen Probes has not had the same capability for analyzing background. It therefore seems fairly certain that some of the older data reported as trapped electrons were actually due to high energy proton contamination. In addition, the situation may also reflect a difference in time periods. The injection of > 1.5-MeV electrons into the inner zone may require extreme magnetic storms while the storms during the Van Allen Probes era have been fairly mild.

This brings up the question of how TID requirements for inner zone missions are affected. As an example the LEO corresponding to the Hubble Space Telescope is examined and presented in Fig. 37. Electron fluence-energy spectra are shown calculated with two models. The first is the AE8 model, which consists of older data from the 1960s to 1970s. The other is AE9/IRENE, which is based on Van Allen Probes data and CRRES data for the inner zone. The only nonzero electron fluxes in AE9/IRENE are due to the CRRES data, which is mainly the result of the severe storm of March 1991. It is seen that the models agree well out to energies of about 1 MeV. Above this, it is not surprising from the above discussion that




Fig. 37. Comparison of the AE8 and AE9/IRENE (version 1.5) models for the LEO of the Hubble Space Telescope.

the AE8 model shows higher fluxes. Analysis of TID behind 2.5 mm of aluminum shielding for the Hubble orbit shows that if AP8/AE8 is used electrons contribute less than 20% of the TID. If AP9/AE9/IRENE is used electrons contribute less than 2% of the TID. The newer model shows inner belt electrons are less significant. Thus, although they present an interesting




Fig. 38. Dose-depth curves for a highly elliptical orbit using two specification methods. The orbit of 1.2×12 earth radii (1274 km perigee \times 70080 km apogee) at a 28.5° inclination includes trapped protons, trapped electrons, and solar protons contributing to TID.

scientific challenge, inner belt electrons are unlikely to drive radiation effect problems except possibly surface effects.

E. Example Environments Including Shielding

1) *Total Ionizing Dose*: TID versus shielding depth curves will be compared using both the traditional margin-based approach and a confidence level based approach that is now possible with the new environment models. A highly elliptical orbit is chosen because it is exposed to all particle populations that contribute significantly to TID – trapped protons, trapped electrons and solar protons. The orbital parameters used were those of the first portion of the NASA Magnetic MultiScale (MMS) mission. Shielding calculations were done for a solid aluminum sphere geometry with dose in silicon calculated at the center of the sphere.

For the margin-based approach, the dose-depth curve is calculated from the mean fluence-energy spectra for trapped particles and solar protons. Energy spectra are obtained from the AP8, AE8 and ESP/PSYCHIC models. Margin is then applied to the dose-depth curve. In this case a margin of times 2 is applied, which is often used by government organizations. This result is shown in Fig. 38. It is compared to the confidence level-based approach in Fig. 38, where the dose-depth curve is calculated from the 95% confidence level fluence-energy spectra obtained from the AP9/AE9/IRENE and ESP/PSYCHIC models. No additional margin is applied to this. It is seen that the results for both approaches agree well out to about 6 mm of aluminum shielding. Beyond this, the difference is primarily due to greater high-energy proton flux levels predicted by AP9/IRENE. A secondary reason is that the newer trapped particle models extend to higher proton and electron energies. For those readers interested in transitioning to the confidence level based approach, dose-depth curves at the 95% confidence level are fairly consistent with using a mean environment and times 2 margin for various orbits.

The confidence level-based TID approach has several advantages over the traditional margin based approach. When convolved with laboratory test data, it allows the device TID failure probability for a given level of shielding to be calculated for the mission [99]. An example of this is shown




Fig. 39. Failure probability for Solid State Devices, Inc., SFT2907A bipolar transistors as a function of shielding level for various orbits [99].

in Fig. 39 for several orbits for bipolar transistors that are used for high-speed, low-power applications. It can be argued this is a better characterization of a device radiation performance in space. It also allows more systematic trades during the design process and is amenable to reliability analyses, which is not possible if only a TID margin is known.

2) *Single Event Upset*: Next SEE environments are considered. The examples presented here are restricted to single event upset (SEU) data and calculations. Fig. 40 shows SEU data from the AFRL SeaStar spacecraft detected on a solid state recorder for more than 4 years [100]. The spacecraft was in a polar LEO. The SEU count per day is shown on the y-axis. There is a slowly varying background of upsets due to trapped protons and GCR. In this case, it is believed most of these SEU were due to trapped protons. Superimposed on this background are sharp increases in the upset rate due to radiation bursts from solar particle events. The largest event spikes were due to the July 14, 2000–July 15, 2000 and November 9, 2000 events. In addition rate spikes due to subsequent smaller events are also seen. Although the environment shown in Fig. 40 is different than what is observed in Fig. 17, note the general similarity in that the effects are due to background radiation that varies slowly with solar cycle superimposed with solar particle events.

Finally, calculated SEU rates are shown for the same highly elliptical orbit considered previously in Fig. 38. In this orbit, the calculations must account for GCR and solar heavy ions. Additionally if the device is sensitive to proton-induced upset, solar protons, and trapped protons must also be considered. SEU rates are shown in Fig. 41 that were calculated for a 4-Gbit NAND flash memory [101]. The sensitive volume was obtained from process reverse engineering and publicly available data. It is seen that increased shielding reduces SEU rates for the worst case solar particle event, the October 1989 event, used in the CREME96 suite of programs. Proton-induced upsets, both those caused by solar protons and trapped protons, can also be reduced with increased shielding. However, the upset rates due to GCR are fairly constant with increased shielding due to their energetic and penetrating nature. Thus, the GCR environment provides a lower limit for the SEU rate that is not practical to reduce significantly. The rates provided




Fig. 40. SEU count per day for a solid state recorder on AFRL’s SeaStar spacecraft in a polar LEO. Data were obtained for more than 4 years beginning in January 1999. Upsets due to the solar particle events of July and November 2000 are identified [100].




Fig. 41. SEU rates calculated for a 4-Gbit NAND flash memory for a worst case solar particle event (CREME96 worst 5 min, worst day and worst week), solar protons (PSYCHIC), trapped protons (AP-8) and GCR during solar minimum and solar maximum as a function of shielding [101].

here for heavy ions do not include fragmentation processes in shielding. Discussion of this is provided in [102] for both SEU and TID.

V. SUMMARY

This paper presented a space climatology timeline ranging from the Big Bang to the present. It began with a description of the early universe including the origin and abundances of particles significant for radiation effects. It continued to

a transition period to modern times when the era of modern space climatology began to emerge due to discoveries of sunspots and the solar activity cycle, along with development of early astronomical methods. The timeline concluded in the modern era that featured the discovery of energetic space radiations and their effects on spacecraft electronics.

ACKNOWLEDGMENT

The author would like to thank P. O’Neill of the Johnson Space Center for providing the calculations and data for Figs. 12, 13, and 15. He would also like to thank C. Stauffer of AS&D, Inc., for assistance with calculations and M. O’Bryan of AS&D, Inc., for assistance with graphics.

REFERENCES

- [1] S. W. Hawking, *A Brief History of Time*. New York, NY, USA: Bantam Books, 1988.
- [2] S. W. Hawking, *A Briefer History of Time*. New York, NY, USA: Bantam Dell, 2005.
- [3] M. Livio, *Brilliant Blunders*. New York, NY, USA: Simon and Schuster, 2013.
- [4] J. C. Mather and J. Boslough, *The Very First Light*. New York, NY, USA: Basic Books, 2008.
- [5] D. Christian, *Origin Story*. New York, NY, USA: Little, Brown & Company, 2018.
- [6] M. Livio, “Hubble’s top ten scientific discoveries,” presentation at the IEEE NSREC, Seattle, WA, USA, Jul. 2005.
- [7] F. Hoyle, “The synthesis of the elements from hydrogen,” *Monthly Notices Roy. Astronom. Soc.*, vol. 106, no. 5, pp. 343–383, Oct. 1946.
- [8] D. Kasen, B. Metzger, J. Barnes, E. Quataert, and E. Ramirez-Ruiz, “Origin of the heavy elements in binary neutron-star mergers from a gravitational-wave event,” *Nature*, vol. 551, pp. 80–84, Nov. 2017, doi: 10.1038/nature24453.
- [9] E. Anders and N. Grevesse, “Abundances of the elements: Meteoritic and solar,” *Geochim. Cosmochim. Acta*, vol. 53, pp. 197–214, Jan. 1989.

- [10] WDC-SILSO. (Sep. 2018). *World Data Center—Sunspot Number and Long-term Solar Observations, Royal Observatory of Belgium, Sunspot Number Catalogue*. [Online]. Available: <http://www.sidc.be/SILSO/>
- [11] B. Rossi, *Cosmic Rays*. New York, NY, USA: McGraw-Hill, 1964.
- [12] S. E. Forbush, "Three unusual cosmic-ray increases possibly due to charged particles from the sun," *Phys. Rev.*, vol. 70, pp. 771–772, Oct. 1946.
- [13] K. F. Galloway, R. L. Pease, R. Schrimpf, and D. W. Emily, "From displacement damage to ELDRS: Fifty years of bipolar transistor radiation effects at the NSREC," *IEEE Trans. Nucl. Sci.*, vol. 60, no. 3, pp. 1731–1739, Jun. 2013.
- [14] R. L. Pease, "A brief history of the NSREC," *IEEE Trans. Nucl. Sci.*, vol. 60, no. 3, pp. 1668–1673, Jun. 2013.
- [15] D. Binder, E. C. Smith, and A. B. Holman, "Satellite anomalies from galactic cosmic rays," *IEEE Trans. Nucl. Sci.*, vol. NS-22, no. 6, pp. 2675–2680, Dec. 1975.
- [16] E. Petersen, R. Koga, M. A. Shoga, J. C. Pickel, and W. E. Price, "The single event revolution," *IEEE Trans. Nucl. Sci.*, vol. 60, no. 3, pp. 1824–1835, Jun. 2013.
- [17] M. A. Xapsos, P. M. O'Neill, and T. P. O'Brien, "Near-Earth space radiation models," *IEEE Trans. Nucl. Sci.*, vol. 60, no. 3, pp. 1691–1705, Jun. 2013.
- [18] S. P. Swordy, "The energy spectra and anisotropies of cosmic rays," *Space Sci. Rev.*, vol. 99, pp. 85–94, Oct. 2001.
- [19] A. I. Mrigakshi, D. Matthia, T. Berger, G. Reitz, and R. F. Wimmer-Schweingruber, "Assessment of galactic cosmic ray models," *J. Geophys. Res.*, vol. 117, p. 8109, Aug. 2012.
- [20] K. Greisen, "End to the cosmic-ray spectrum?" *Phys. Rev. Lett.*, vol. 16, pp. 748–750, Apr. 1966.
- [21] J. H. Adams, Jr., R. Silberberg, and C. H. Tsao, "Cosmic ray effects on microelectronics. Part 1: The near-Earth particle environment," Naval Res. Lab., Washington, DC, USA, NRL Memorandum Rep. 4506, Aug. 1981.
- [22] J. H. Adams, Jr., "Cosmic ray effects on microelectronics. Part 4," Naval Res. Lab., Washington, DC, USA, NRL Memorandum Rep. 5901, Dec. 1987.
- [23] A. J. Davis *et al.*, "Solar minimum spectra of galactic cosmic rays and their implications for models of the near-Earth radiation environment," *J. Geophys. Res.*, vol. 106, no. A12, pp. 29979–29987, Dec. 2001.
- [24] N. V. Kuznetsov, H. Popova, and M. I. Panasyuk, "Empirical model of long-time variations of galactic cosmic ray particle fluxes," *J. Geophys. Res. Space Phys.*, vol. 122, pp. 1463–1472, Feb. 2017.
- [25] F. Lei, A. Hands, S. Clucas, C. Dyer, and P. Truscott, "Improvements to and validations of the QinetiQ atmospheric radiation model (QARM)," *IEEE Trans. Nucl. Sci.*, vol. 53, no. 4, pp. 1851–1858, Aug. 2006.
- [26] D. Matthia, T. Berger, A. I. Mrigakshi, and G. Reitz, "A ready-to-use galactic cosmic ray model," *Adv. Space Res.*, vol. 51, no. 3, pp. 329–338, 2013.
- [27] R. A. Nymmik, M. I. Panasyuk, and A. A. Suslov, "Galactic cosmic ray flux simulation and prediction," *Adv. Space Res.*, vol. 17, no. 2, pp. 19–30, 1996.
- [28] A. J. Tylka *et al.*, "CREME96: A revision of the cosmic ray effects on micro-electronics code," *IEEE Trans. Nucl. Sci.*, vol. 44, no. 6, pp. 2150–2160, Dec. 1997.
- [29] P. M. O'Neill, "Badhwar–O'Neill 2010 galactic cosmic ray flux model—Revised," *IEEE Trans. Nucl. Sci.*, vol. 57, no. 6, pp. 3148–3153, Dec. 2010.
- [30] P. M. O'Neill, S. Golge, and T. C. Slaba, "Badhwar–O'Neill 2014 galactic cosmic ray flux model description," NASA Johnson Space Center, Houston, TX, USA, Tech. Rep. NASA/TP-2015-218569, Mar. 2015.
- [31] E. L. Petersen, J. C. Pickel, J. H. Adams, and E. C. Smith, "Rate prediction for single event effects—A critique," *IEEE Trans. Nucl. Sci.*, vol. 39, no. 6, pp. 1577–1599, Dec. 1992.
- [32] R. A. Reed *et al.*, "Impact of ion energy and species on single event effects analysis," *IEEE Trans. Nucl. Sci.*, vol. 54, no. 6, pp. 2312–2321, Dec. 2007.
- [33] D. V. Reames, "Particle acceleration at the sun and in the heliosphere," *Space Sci. Rev.*, vol. 90, pp. 413–491, Oct. 1999.
- [34] J. L. Barth, "Modeling space radiation environments," in *Proc. IEEE NSREC*, Piscataway, NJ, USA, Jul. 1997, pp. I-1–I-82.
- [35] M. A. Xapsos, C. Stauffer, J. L. Barth, and E. A. Burke, "Solar particle events and self-organized criticality: Are deterministic predictions of events possible?" *IEEE Trans. Nucl. Sci.*, vol. 53, no. 4, pp. 1839–1843, Aug. 2006.
- [36] J. H. King, "Solar proton fluences for 1977–1983 space missions," *J. Spacecraft*, vol. 11, pp. 401–408, Jun. 1974.
- [37] E. G. Stassinopoulos and J. H. King, "Empirical solar proton model for orbiting spacecraft applications," *IEEE Trans. Aerosp. Electron. Syst.*, vol. AES-10, no. 4, pp. 442–450, Jul. 1974.
- [38] J. Feynman, T. P. Armstrong, L. Dao-Gibner, and S. Silverman, "New interplanetary proton fluence model," *J. Spacecraft*, vol. 27, pp. 403–410, Jul./Aug. 1990.
- [39] J. Feynman, G. Spitale, J. Wang, and S. Gabriel, "Interplanetary proton fluence model: JPL 1991," *J. Geophys. Res.*, vol. 98, pp. 13281–13294, Aug. 1993.
- [40] R. A. Nymmik, "Probabilistic model for fluences and peak fluxes of solar energetic particles," *Radiat. Meas.*, vol. 30, pp. 287–296, Jun. 1999.
- [41] M. A. Xapsos, G. P. Summers, J. L. Barth, E. G. Stassinopoulos, and E. A. Burke, "Probability model for worst case solar proton event fluences," *IEEE Trans. Nucl. Sci.*, vol. 46, no. 6, pp. 1481–1485, Dec. 1999.
- [42] M. A. Xapsos, G. P. Summers, J. L. Barth, E. G. Stassinopoulos, and E. A. Burke, "Probability model for cumulative solar proton event fluences," *IEEE Trans. Nucl. Sci.*, vol. 47, no. 3, pp. 486–490, Jun. 2000.
- [43] P. T. A. Jiggins, S. B. Gabriel, D. Heynderickx, N. Crosby, A. Glover, and A. Hilgers, "ESA SEP-EM project: Peak flux and fluence model," *IEEE Trans. Nucl. Sci.*, vol. 59, no. 4, pp. 1066–1077, Aug. 2012.
- [44] P. Jiggins, D. Heynderickx, I. Sandberg, P. Truscott, O. Raukunen, and R. Vainio, "Updated model of the solar energetic proton environment in space," *J. Space Weather Space Clim.*, vol. 8, May 2018, Art. no. A31.
- [45] Z. D. Robinson, J. H. Adams, M. A. Xapsos, and C. A. Stauffer, "Database of episode-integrated solar energetic proton fluences," *J. Space Weather Space Clim.*, vol. 8, p. A24, Jan. 2018.
- [46] J. H. Adams, Jr., W. F. Dietrich, and M. A. Xapsos, "Probabilistic solar energetic particle models," in *Proc. 32nd Int. Cosmic Ray Conf.*, Beijing, China, vol. 10, 2011, pp. 20–23.
- [47] M. A. Xapsos, "Modeling the space radiation environment," in *Proc. IEEE NSREC*, Piscataway, NJ, USA, Jul. 2006, pp. I-1–I-62.
- [48] I. Jun, R. T. Swimm, A. Ruzmaikin, J. Feynman, A. J. Tylka, and W. F. Dietrich, "Statistics of solar energetic particle events: Fluences, durations, and time intervals," *Adv. Space Res.*, vol. 40, no. 3, pp. 304–312, 2007.
- [49] O. Raukunen *et al.*, "Two solar proton fluence models based on ground level enhancement observations," *J. Space Weather Space Clim.*, vol. 8, Jan. 2018, Art. no. A04.
- [50] A. J. Tylka and W. F. Dietrich, "A new and comprehensive analysis of proton spectra in ground-level enhanced (GLE) solar particle events," in *Proc. 31st Int. Cosmic Ray Conf.*, Łódź, Poland, 2009, pp. 7–15.
- [51] M. A. Xapsos, C. Stauffer, G. B. Gee, J. L. Barth, E. G. Stassinopoulos, and R. E. McGuire, "Model for solar proton risk assessment," *IEEE Trans. Nucl. Sci.*, vol. 51, no. 6, pp. 3394–3398, Dec. 2004.
- [52] P. Truscott *et al.*, "Methods for and the influence of uncertainty propagation in the solar energetic particle environment modelling (SEP-EM) system," presentation at the IEEE NSREC, Kona, HI, USA, Jul. 2018.
- [53] M. A. Xapsos, C. Stauffer, T. Jordan, J. L. Barth, and R. A. Mewaldt, "Model for cumulative solar heavy ion energy and linear energy transfer spectra," *IEEE Trans. Nucl. Sci.*, vol. 54, no. 6, pp. 1985–1989, Dec. 2007.
- [54] B. J. Anderson and R. E. Smith, "Natural orbital environment definition guidelines for use in aerospace vehicle development," NASA Marshall Space Flight Center, Huntsville, AL, USA, NASA Tech. Memorandum 4527, Jun. 1994.
- [55] R. A. Nymmik, "Models describing solar cosmic ray events," *Radiat. Meas.*, vol. 26, pp. 417–420, May 1996.
- [56] E. G. Stassinopoulos, G. J. Brucker, D. W. Nakamura, C. A. Stauffer, G. B. Gee, and J. L. Barth, "Solar flare proton evaluation at geostationary orbits for engineering applications," *IEEE Trans. Nucl. Sci.*, vol. 43, no. 2, pp. 369–382, Apr. 1996.
- [57] K. G. McCracken, G. A. M. Dreschhoff, E. J. Zeller, D. F. Smart, and M. A. Shea, "Solar cosmic ray events for the period 1561–1994: 1. Identification in polar ice," *J. Geophys. Res.*, vol. 106, pp. 21585–21598, Oct. 2001.
- [58] E. W. Wolff *et al.*, "The Carrington event not observed in most ice core nitrate records," *Geophys. Res. Lett.*, vol. 39, no. 8, p. L08503, Apr. 2012.

- [59] C. S. Dyer, K. Hunter, S. Clucas, D. Rodgers, A. Campbell, and S. Buchner, "Observation of solar particle events from CREDO and MPTB during the current solar maximum," *IEEE Trans. Nucl. Sci.*, vol. 49, no. 6, pp. 2771–2775, Dec. 2002.
- [60] M. A. Xapsos, G. P. Summers, and E. A. Burke, "Probability model for peak fluxes of solar proton events," *IEEE Trans. Nucl. Sci.*, vol. 45, no. 6, pp. 2948–2953, Dec. 1998.
- [61] E. Gumbel, *Statistics of Extremes*. New York, NY, USA: Columbia Univ. Press, 1958.
- [62] A. H.-S. Ang and W. H. Tang, *Probability Concepts in Engineering Planning and Design*, vol. 2, New York, NY, USA: Wiley, 1975.
- [63] E. Castillo, *Extreme Value Theory in Engineering*. Boston, MA, USA: Academic, 1988.
- [64] P. J. McNulty, L. Z. Scheick, D. R. Roth, M. G. Davis, and M. R. S. Tortora, "First failure predictions for EPROMs of the type flown on the MPTB satellite," *IEEE Trans. Nucl. Sci.*, vol. 47, no. 6, pp. 2237–2243, Dec. 2000.
- [65] P. J. Vail and E. A. Burke, "Fundamental limits imposed by gamma dose fluctuations in scaled MOS gate insulators," *IEEE Trans. Nucl. Sci.*, vol. NS-31, no. 6, pp. 1411–1416, Dec. 1984.
- [66] M. A. Xapsos, "Hard error dose distributions of gate oxide arrays in the laboratory and space environments," *IEEE Trans. Nucl. Sci.*, vol. 43, no. 6, pp. 3139–3144, Dec. 1996.
- [67] E. A. Burke *et al.*, "Gamma induced dose fluctuations in a charge injection device," *IEEE Trans. Nucl. Sci.*, vol. NS-35, no. 6, pp. 1302–1306, Dec. 1988.
- [68] P. W. Marshall, C. J. Dale, E. A. Burke, G. P. Summers, and G. E. Bender, "Displacement damage extremes in silicon depletion regions," *IEEE Trans. Nucl. Sci.*, vol. 36, no. 6, pp. 1831–1839, Dec. 1989.
- [69] E. T. Jaynes, "Information theory and statistical mechanics," *Phys. Rev.*, vol. 106, no. 4, pp. 620–630, 1957.
- [70] J. N. Kapur, *Maximum Entropy Models in Science and Engineering*. New York, NY, USA: Wiley, 1989.
- [71] R. A. Nymmik, "Improved environment radiation models," *Adv. Space Res.*, vol. 40, no. 3, pp. 313–320, 2007.
- [72] M. A. Xapsos, G. P. Summers, and E. A. Burke, "Extreme value analysis of solar energetic proton peak fluxes," *Sol. Phys.*, vol. 183, pp. 157–164, Nov. 1998.
- [73] S. Bourdarie and M. Xapsos, "The near-Earth space radiation environment," *IEEE Trans. Nucl. Sci.*, vol. 55, no. 4, pp. 1810–1832, Aug. 2008.
- [74] P. Jiggins *et al.*, "The solar accumulated and peak proton and heavy ion radiation environment (SAPPHIRE) model," *IEEE Trans. Nucl. Sci.*, vol. 65, no. 2, pp. 698–711, Feb. 2018.
- [75] B. Taylor, G. Vacanti, E. Maddox, and C. I. Underwood, "The interplanetary electron model (IEM)," *IEEE Trans. Nucl. Sci.*, vol. 58, no. 6, pp. 2785–2792, Dec. 2011.
- [76] C. E. McIlwain, "Coordinates for mapping the distribution of magnetically trapped particles," *J. Geophys. Res.*, vol. 66, no. 11, pp. 3681–3691, Nov. 1961.
- [77] M. Walt, *Introduction to Geomagnetically Trapped Radiation*. Cambridge, MA, USA: Cambridge Univ. Press, 1994.
- [78] S. L. Huston and K. A. Pfützer, "A new model for the low altitude trapped proton environment," *IEEE Trans. Nucl. Sci.*, vol. 45, no. 6, pp. 2972–2978, Dec. 1998.
- [79] W. R. Johnston, T. P. O'Brien, S. L. Huston, T. B. Guild, and G. P. Ginot, "Recent updates to the AE9/AP9/SPM radiation belt and space plasma specification model," *IEEE Trans. Nucl. Sci.*, vol. 62, pp. 2760–2766, Dec. 2015.
- [80] M. S. Gussenhoven, E. G. Mullen, and D. H. Brautigam, "Improved understanding of the Earth's radiation belts from the CRRES satellite," *IEEE Trans. Nucl. Sci.*, vol. 43, no. 2, pp. 353–368, Apr. 1996.
- [81] D. M. Sawyer and J. I. Vette, "AP-8 trapped proton environment for solar maximum and solar minimum," NASA Goddard Space Flight Center, Greenbelt, MD, USA, Tech. Rep. NSSDC/WDC-A-R&S, Dec. 1976, p. 76.
- [82] J. I. Vette, "The NASA/National Space Science Data Center trapped radiation environment model program, 1964–1991," NASA Goddard Space Flight Center, Nat. Space Sci. Data Center, Greenbelt, MD, USA, Tech. Rep. NSSDC/WDC-A-R/S-91-29, Nov. 1991.
- [83] E. J. Daly, J. Lemaire, D. Heynderickx, and D. J. Rodgers, "Problems with models of the radiation belts," *IEEE Trans. Nucl. Sci.*, vol. 43, no. 2, pp. 403–414, Apr. 1996.
- [84] D. Heynderickx, M. Kruglanski, V. Pierrard, J. Lemaire, M. D. Looper, and J. B. Blake, "A low altitude trapped proton model for solar minimum conditions based on SAMPEX/PET data," *IEEE Trans. Nucl. Sci.*, vol. 46, no. 6, pp. 1475–1480, Dec. 1999.
- [85] S. L. Huston, "Space environments and effects: Trapped proton model," Boeing Co., Huntington Beach, CA, USA, Final Rep. NAS8-98218, Jan. 2002.
- [86] D. Boscher, A. Sicard-Piet, D. Lazaro, T. Cayton, and G. Rolland, "A new proton model for low altitude high energy specification," *IEEE Trans. Nucl. Sci.*, vol. 61, no. 6, pp. 3401–3407, Dec. 2014.
- [87] J.-M. Lauenstein and J. L. Barth, "Radiation belt modeling for spacecraft design: Model comparisons for common orbits," in *Proc. IEEE Radiation Effects Data Workshop*, Piscataway, NJ, USA, 2005, pp. 102–109.
- [88] G. P. Ginot *et al.*, "AE9, AP9 and SPM: New models for specifying the trapped energetic particle and space plasma environment," *Space Sci. Rev.*, vol. 179, nos. 1–4, pp. 579–615, Nov. 2013.
- [89] A. Sicard-Piet *et al.*, "GREEN: The new global radiation Earth environment model (beta version)," *Ann. Geophys.*, vol. 36, pp. 953–967, Jul. 2018.
- [90] J. I. Vette, "The AE-8 trapped electron model environment," NASA Goddard Space Flight Center, Greenbelt, MD, USA, Tech. Rep. NSSDC/WDC-A-RS-91-24, Nov. 1991.
- [91] W. D. Pesnell, "Fluxes of relativistic electrons in low-Earth orbit during the decline of solar cycle 22," *IEEE Trans. Nucl. Sci.*, vol. 48, no. 6, pp. 2016–2021, Dec. 2001.
- [92] G. L. Wrenn, D. J. Rodgers, and P. Buehler, "Modeling the outer belt enhancements of penetrating electrons," *J. Spacecraft Rockets*, vol. 37, no. 3, pp. 408–415, May/June 2000.
- [93] H. C. Koons, "Statistical analysis of extreme values in space science," *J. Geophys. Res.*, vol. 106, no. A6, pp. 10915–10921, Jun. 2001.
- [94] D. H. Brautigam, K. P. Ray, G. P. Ginot, and D. Madden, "Specification of the radiation belt slot region: Comparison of the NASA AE8 model with TSX5/CEASE data," *IEEE Trans. Nucl. Sci.*, vol. 51, no. 6, pp. 3375–3380, Dec. 2004.
- [95] D. H. Brautigam, M. S. Gussenhoven, and E. G. Mullen, "Quasi-static model of outer zone electrons," *IEEE Trans. Nucl. Sci.*, vol. 39, no. 6, pp. 1797–1803, Dec. 1992.
- [96] A. L. Vampola, "The ESA outer zone electron model update," in *Proc. Environ. Modelling Space-Based Appl., Symp. (ESA SP)*, W. Burke and T.-D. Guyenne, Eds. Noordwijk, The Netherlands: ESTEC, Sep. 1996, pp. 18–20.
- [97] A. Sicard-Piet *et al.*, "A new international geostationary electron model: IGE-2006, from 1 keV to 5.2 MeV," *Space Weather*, vol. 6, no. 7, pp. 1–13, Jul. 2008.
- [98] S. G. Claudepierre *et al.*, "The hidden dynamics of relativistic electrons (0.7–1.5 MeV) in the inner zone and slot region," *J. Geophys. Res.*, *Space Phys.*, vol. 122, no. 3, pp. 3127–3144, Mar. 2017.
- [99] M. A. Xapsos *et al.*, "Inclusion of radiation environment variability in total dose hardness assurance methodology," *IEEE Trans. Nucl. Sci.*, vol. 64, no. 1, pp. 325–331, Jan. 2017.
- [100] C. Poivrey *et al.*, "Single event upset (SEU) study of SeaStar, and the MAP anomaly," presented at the Single Event Effects Symp., Los Angeles, CA, USA, Apr. 2002.
- [101] J. A. Pellish *et al.*, "Impact of spacecraft shielding on direct ionization soft error rates for sub-130 nm technologies," *IEEE Trans. Nucl. Sci.*, vol. 57, pp. 3183–3189, Dec. 2010.
- [102] P. M. O'Niell, "Ionizing radiation environment inside spacecraft," in *Proc. IEEE NSREC*, Piscataway, NJ, USA, Jul. 2015, pp. 1-1–1-55.